

Special Advisory LAB-LINK

New and updated laboratory testing information

April 2023

New Genomics Test — NOW AVAILABLE!

al Screening NIPT
Noninvasive Prenatal Screening
NIPT
81420
Noninvasive prenatal screening for trisomy 13, 18, and 21 and sex chromosome aneuploidies in singleton and twin pregnancies.
Cell Free DNA; Noninvasive Prenatal Aneuploidy Screen; cfDNA
Next Generation Sequencing of Cell-Free DNA
Routine, at least twice per week
4-6 days; report available in EPIC
rs
7-10 mL whole blood, collect two tubes
Streck Cell-Free BCT, collect two tubes
Draw blood using Vacutainer system (butterfly not appropriate); invert tube 10 times.
After samples are received, store between 2-8 degrees Celsius before ready to proceed.
Noninvasive prenatal screening for trisomy 13, 18, and 21 and sex chromosome aneuploidies. Recommended by ACOG for any pregnant woman, particularly those with a high-risk pregnancy.
Disomy for chromosomes 13, 18, and 21 and XX or XY sex chromosomes.
Time between collection and initial plasma isolation should not exceed 5 days.

For technical questions related to this test, please call HNL Lab Medicine's genetic testing department, at 484-244-2900.

For general inquiries, Customer Care is available to assist at any time by calling 877-402-4221.