March 28, 2025

New Methodology

Compliance Oral Fluid Drug Screening

Description of Change

Starting **March 31, 2025**, screening tests for Oral Fluid testing will be updated with new methodology and greater sensitivity. As a result, the following tests and cutoff concentrations will be impacted. The Current Single Order test codes are being replaced with the New Single Order Test Codes.

Analytes (Test Code)	Current Single Order Test Code	NEW Single Order Test Code	Current Cutoff CONC (ng/mL)	NEW Cutoff CONC (ng/mL)
COMPLIANCE DRUG MONITORING DM BASIC PROFILE, OF (OCBP)				
AMPHETAMINES	AMPHO	**APHTO**	150	30
METHAMPHETAMINE	AMAPO	**AMAO**	120	30
BARBITURATES	BABTO	**BARBO**	60	60
BENZODIAZEPINES	BNDZO	**BNDO**	3	1.2
CANNABINOIDS	CANNO	**CANO**	3	3
COCAINE METABOLITE	COCMO	**COCCO**	15	6
OPIATES	OPISO	**OPIO**	30	6
OXYCODONE	OXYSO	**OCODO**	30	3
METHADONE	MTHAO	**MTHO**	15	6
PHENCYCLIDINE - PCP	PHCO	**PNCYO**	3	3
BUPRENORPHINE	BPRNO	**BUPNO**	3	0.6
FENTANYL not part of OCBP (order separately)			N/A	0.6

Effective Date	**Monday, March 31, 2025**
Test Name	Drug Screen, Qualitative, Oral Fluid
Test Code	OCBP
CPT Code	80307
Methodology	Current: Enzyme Immunoassay (EIA) **NEW: Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)**
Testing Schedule	Routine, daily
Report Availability	1 - 5 days (for confirmation of positive screening results or if reflex testing is required)

Minimum Volume	1 mL
Container	Intercept®i2he™ Oral Fluid Collection Device Specimens received in any container other than the Intercept®i2he™ Oral Fluid Collection Device will be rejected.
Collection Instructions	1. Confirm the donor has not had anything in their mouth for 10 minutes prior to providing the oral fluid sample. If the donor has had anything in their mouth within the last 10 minutes, wait 10 minutes prior to the collection process. 2. Instruct the donor to remove the device from the side of the pouch labeled #1 by grabbing the plastic shaft of the collection pad. Ensure the donor does not touch the pad. 3. Instruct the donor to place the collection pad under the tongue with white side of the stick facing upward. Instruct the donor not to bite down on the device but to grip between the lips. 4. Instruct the donor to keep the pad under the tongue until the indicator turns blue. Should the collection time reach 15 minutes and the indicator has NOT turned blue, stop the collection and proceed. 5. Instruct the donor to remove the device from their mouth and slowly place the collection device into the vial (side of pouch labeled #2). 6. Cap the vial.
Reference Range	• Interpretation and concentration of drug/metabolite in oral fluid is dependent on dose, time of dose, and metabolic rate. • Reported results are corrected for the collection device dilution.

Additional Information

If you have additional questions regarding this test, please contact Technical Support between the hours of 8 a.m. and 4:30 p.m. For general questions, Customer Care is available to assist at any time.

Technical Support/Customer Care

Call 877-402-4221